

The Effects of Low-Shear Simulated Microgravity on Soil Bacterium *Pantoea* *agglomerans*

Steven Russell

Summer 2019 NIFS Intern

NASA John F. Kennedy Space Center, FL 32899

University of North Dakota, Grand Forks, ND 58202

Objectives



- Determine growth rate of *Pantoea agglomerans* in simulated microgravity and normal gravity conditions.
- Determine antibiotic susceptibility of *Pantoea agglomerans* after exposure to simulated microgravity and normal gravity conditions.
- Have Fun 😊

Materials

- Rotary Cell Culture System (RCCS)
- High Aspect-Ratio Vessel (HARV)
- Strain of *Pantoea agglomerans* (*ISS Isolate*)
- Spectrophotometer
- Trypticase Soy Broth (TSB) and Agar (TSA) Plates
- Mueller-Hinton Agar Plates
- Antibiotic Infused Disks
- GraphPad Prism (Statistical Analysis)
- High Spirit and a Good Attitude ☺

Panotea agglomerans

- *P. agglomerans*
- Gram-negative
- Rod-Shaped
- Motile – peritrichous
- Enterobacteriaceae Family (*E. coli*, *Klebsiella*, *Salmonella*, *Shigella*, and *Yersinia pestis*)



Low- Shear Simulated Microgravity (LSSMG)

- RCCS
- HARV
- Low Shear – no bubbles in vessel once filled
- Constant rotation rate
- Vertical Rotation (Fig.1) = Simulated Microgravity
- Horizontal No Rotation (Fig.2) = Normal Gravity




Fig.1





Fig. 2







Relevance


- Plant Growth Promoter (Dutkiewicz et. al; Mirsha et. Al)
- AND Plant Pathogen (Cruz et. al, Dutkiewicz et. al)
- Opportunistic Pathogen (Büyükcama et. al, Cruz et. al)
- Immunosuppressed Astronauts (Mann et. al)
- Microbial growth in microgravity (Nickerson et. al)

Hypotheses


- H_{1A} – *P. agglomerans* growth rate will increase in LSSMG compared to normal gravity.
- H_{2A} – *P. agglomerans* growth rate will decrease in LSSMG compared to normal gravity.
- H_{0A} – There will be no change in the growth rate of *P. agglomerans* in LSSMG compared to normal gravity.
- H_{1B} – *P. agglomerans* will become less susceptible to chosen antibiotics after exposure LSSMG compared to normal gravity.
- H_{0B} – *P. agglomerans* will remain susceptible to chosen antibiotics after exposure to LSSMG compared to normal gravity.

Experimental Design – Exp. 1


- Determine growth rate in a shaking flask
- Determine growth rate in LSSMG and normal gravity
 - Experimental Rotated HARVS vertically
 - Horizontal Control HARVS
 - Shaking flask as ground control
 - Removed sample
 - Read optical density (OD)
 - Plated to confirm starting concentration




Exp. 1 Trial 1




Exp. 1 Trial 2 Results



Exp. 1 Trial 2 Results – Cont.




Exp. 1 Trial 1 & 2 Compared



Experimental Design – Exp. 2


- “Time Point” Experiment
 - Chose 4 time points
 - 2 vessels per time point
 - Experimental (vertical) and control (horizontal) – 8 vessels total
 - Removed 2 vessels at each time point
 - Harvested the entire vessel – OD, plated for counts and for antibiotic test
 - Also ran a shaking culture






Exp. 2 Results –Growth Curve

Experimental v Shaking Exp.2

—●— LSSMG —●— Control —●— Shaking




Exp. 2 Results – Counts



Exp. 2 – Cont.

- Antibiotic Test
 - Kirby-Bauer Antibiotic Susceptibility Test
 - Used to determine antibiotic susceptibility of bacteria
 - Measuring Zone of Inhibition (ZOI)
 - Mueller-Hinton Agar is used.
- Antibiotics
 - Ciprofloxacin (5µg)
 - Tetracycline (30µg)
 - Chloramphenicol (30µg)
- Tested *E. coli* ATC25922 for quality control



Exp. 2 Results – Antibiotics (ZOI)

Antibiotic	Susceptibility	Intermediate	Resistant
Tetracycline (30µg)	≥15mm	12-14mm	≤11mm
Ciprofloxacin (5µg)	≥21mm	16-20mm	≤15mm
Chloramphenicol (30µg)	≥18mm	13-17mm	≤12mm

*

*Based on standards set by the CLSI MS100 for the *Enterobacteriaceae* family.

***P. agglomerans* ZOI Average**

- Control
 - Tetracycline: 22.4mm= S
 - Ciprofloxacin: 34.9mm= S
 - Chloramphenicol: 24.3mm= S
- LSSMG
 - Tetracycline: 21.1mm= S
 - Ciprofloxacin: 34.4mm= S
 - Chloramphenicol: 24.4mm= S
- Shaking Ground Control
 - Tetracycline: 22.4mm= S
 - Ciprofloxacin: 35.4mm= S
 - Chloramphenicol: 24.9mm= S

Conclusions

- So far:
 - Growth Curves: H_{1A} looks promising (Hypothesis 1 is accepted)
 - Antibiotic Resistance: H_{0B} looks likely (Null Hypothesis is accepted)
- Some significant differences in the growth rate in LSSMG compared to the control during 1hr intervals and media addition.
- No significant change in growth rate in LSSMG compared to the control when ran continuously, chose time points and no media addition
- No change in antibiotic susceptibility.
- Tale of two experiments.

Future Work

- Repeat, Repeat and Repeat again
- Continuously run bacterial culture in RCCS
 - Antibiotic resistance may not have had time to develop
- Addition of an antibiotic resistant microbial species.
 - Test in LSSMG if there is gene transfer of antibiotic resistance gene to *P. agglomerans*
- Addition of Media vs Harvesting entire Vessel
- Kirby-Bauer on Addition of Media Experiments

Acknowledgements

- Anna Maria Ruby, NASA KSC
 - Dr. Ye Zhang, NASA KSC
- Mary Hummerick, AECOM LASSO
- Dr. Christina Khodadad, AECOM LASSO
 - Jeff Richards, AECOM LASSO
- North Dakota Space Grant Consortium, University of North Dakota
 - Joey Emhof, NIFS
- The Rest of the 2019 NIF Summer Interns in the “Petri Dish”

Questions?

References

- Büyükcama, A., Tuncerb, Ö., Gürb, D., Sancakb, B., Ceyhana, M., Cengiza, A.B., Kara, A., “Clinical and microbiological characteristics of Pantoea agglomerans infection in children,” *Journal of Infection and Public Health*, Vol. 11, 2018, pp. 304-309.
- Cruz, A. T., Cazacu, A. C., Allen, C. H., “*Pantoea agglomerans*, a Plant Pathogen Causing Human Disease,” *Journal of Clinical Microbiology*, Vol. 45, No. 6, 2007, pp. 1989-1992.
- Dutkiewicz, J., Mackiewicz, B., Lemieszek, M. K., Golec, M., Milanowski, J., “Pantoea agglomerans: a mysterious bacterium of evil and good. Part III. Deleterious effects: infections of humans, animals and plants,” *Annals of Agriculture and Environmental Science*, Vol. 23, No. 2, 2016, pp. 197-205.
- Mann V, Sundaresan A, Mehta S. K., Crucian B, Doursout M.F., Devakottai S., Effects of microgravity and other space stressors in immunosuppression and viral reactivation with potential nervous system involvement. *Neurology India* Vol. 67 No. 8, 2019, pp. 198-203.
- Mirsha, A., Chauhan, S. P., Chaudhry, V., Tripathi, M., Nautiyal, C. S., “Rhizosphere competent Pantoea agglomerans enhances maize (*Zea mays*) and chickpea (*Cicer arietinum* L.) growth, without altering the rhizosphere functional diversity,” *Antonie van Leeuwenhoek*, Vol. 100, No. 3, 2011, pp. 405-413.
- Nickerson, C., Ott, M., Wilson, J.W., Ramamurthy, R., Pierson, D.L., “Microbial Responses to Microgravity and Other Low-Shear Environments,” *Microbiology and Molecular Biology Reviews*, Vol. 68, No. 2, 2004, pp. 345-361.